

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>7</sup>:

C12N 5/00

A2

(11) International Publication Number:

WO 00/70021

(43) International Publication Date: 23 November 2000 (23.11.00)

(21) International Application Number: PCT/IL00/00270

(22) International Filing Date: 12 May 2000 (12.05.00)

(30) Priority Data:

129966

14 May 1999 (14.05.99)

IL

(71) Applicant (for all designated States except US): YISSUM RESEARCH DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; P.O. Box 4279, 91042 Jerusalem (IL).

(71)(72) Applicant and Inventor: ITS KOVITZ-ELDOR, Joseph [IL/IL]; Sherit Haplayta St. 42, Danya B., 34987 Haifa (IL).

(72) Inventor; and



(75) Inventor/Applicant (for US only): BENVENISTY, Nissim [IL/IL]; Hertzl St. 83, Beit-Hakerem, 96344 Jerusalem (IL).

(74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel Aviv (IL).

(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published***Without international search report and to be republished upon receipt of that report.*

(54) Title: DIFFERENTIATED HUMAN EMBRYOID CELLS AND A METHOD FOR PRODUCING THEM

**A****B****(57) Abstract**

A process for obtaining human derived embryoid bodies (hEB). Human embryonic stem cells are incubated *in vitro* in a liquid growth medium under conditions in which the cells undergo differentiation, but do not adhere to the walls of the container. The invention also provides hEBs obtained by the process.

**BEST AVAILABLE COPY**





## DIFFERENTIATED HUMAN EMBRYOID CELLS AND A METHOD FOR PRODUCING THEM

### FIELD OF THE INVENTION

The present invention is in the field of cell biology and more specifically relates to methods of *in vitro* differentiation.

### BACKGROUND OF THE INVENTION

Embryonic stem (ES) cells are derived from totipotent cells in an embryo. Murine ES cells have been shown to be pluripotent cells capable *in vitro* of terminal differentiation into cells of the mesoderm, ectoderm and endoderm lineages (Robertson, 1987; Dushnik-Levinson and Benvenisty, 1995). The pluripotency of murine embryonic stem cells has been established on the basis of three criteria:

- (1) When undifferentiated murine ES cells are injected into the cavity of a blastocyst and the blastocysts are subsequently implanted into pseudo-pregnant mice, chimeric mice develop. The injected ES cells contribute to all cell types, including the germ layer. Thus, in the next generation, mice with the genotype of the ES cells are born (Capecchi, 1989; Rossant and Joyner, 1989).
- (2) When murine ES cells are injected subcutaneously into syngeneic mice, teratoma tumors develop. These tumors comprise cells of all three embryonic origins (endoderm, ectoderm and mesoderm) (Wobus *et al.*, 1984).

- (3) When murine ES cells are allowed to aggregate *in vitro* so as to form embryoid bodies (EBs), the cells differentiate in the EBs into various cell types (Robertson, 1987).

During maturation of murine ES cells *in vitro*, in addition to morphological changes, a cell may acquire a molecular marker characteristic of a differentiated cell type, such as  $\xi$ -globin (a marker of hematopoietic cells) (Wiles and Keller, 1991; Lindenbaum and Grosveld, 1990), neurofilament-68kd protein (a marker of neuronal cells) (Bain *et al.*, 1995; Levinson-Dushnik and Benvenisty, 1997), and albumin (a marker of hepatic cells) (Levinson-Dushnik and Benvenisty, 1997).

ES cell lines have also been established from primates such as the common marmoset (Thomson, 1996)), and the rhesus monkey (Thomson, 1995)). However, EB formation by marmoset ES cells is inconsistent and asynchronous, and differentiation of the rhesus ES cells is disorganized and vesicular structures do not form (Thomson, 1998a).

Human ES cell lines have been established derived from human embryos produced by *in vitro* fertilization (Thomson, 1998b). The embryos were cultured to the blastocyst stage, and inner cell masses comprising ES cells were isolated and cultured. These human ES cells are only known to be capable of differentiating when in teratomas (Thomson, 1998b). When cultured *in vitro*, the human ES cells have normal karyotypes, express telomerase activity, and proliferate. However, the inability to produce EBs from non-human primate ES cells lead to the belief that EBs could also not be formed from human ES cells (Thomson, 1998a).

## SUMMARY OF THE INVENTION

The present invention is based on the finding that, contrary to the aforementioned belief that human ES cells do not form EBs, under appropriate conditions, human embryoid bodies (hEBs) may be obtained *in vitro*, from human ES (hES) cells suspended in a liquid medium. These hEBs contain mesoderm, ectoderm and endoderm cell lineages and may be used as a source of cells of the different lineages, e.g. for transplantation or inoculation into human recipients. By

incubating the hEBs, these basic cell lineages can differentiate into a wide variety of different cell types, e.g. cells having characteristics of cardiac cells, neural stem cells, and others.

The hEBs in accordance with the invention may be used as a source of cells  
5 for use in transplantation or inoculation into a human recipient, in order to treat various diseases or disorders, to assist in tissue repair, to substitute for degenerated tissue.

Thus, in accordance with one aspect, the present invention provides a human embryoid body (hEB).

10 The hEB, in accordance with the invention, is preferably obtained by *in vitro* culturing of hES cells in a vessel under conditions in which the cells or aggregates thereof do not adhere to the vessel walls.

In accordance with another aspect, the invention provides a process for obtaining at least one human-derived embryoid body (hEB), comprising:

- 15 (a) providing human embryonic stem (hES) cells;  
(b) growing the hES cells *in vitro* in a vessel under conditions in which said cells undergo differentiation and the cells or aggregates thereof do not adhere to the vessel wall; and  
(c) incubating for a time sufficient to develop hEBs from said cells.

20 The conditions whereby the hES cells do not adhere to the vessel wall include culturing the cells in a vessel having walls made of a material to which the cells or aggregates thereof are incapable of adhering, adding one or more factors into the medium which prevent adherence of the cells to the vessel walls etc. Conditions whereby the hES undergo differentiation include the absence of  
25 inhibitors of differentiation such as leukemia inhibitory factor and fibroblast growth factor.

The invention also provides a process for preparing cells of a defined cell lineage comprising:

- (a) providing human embryonic stem (hES) cells;

(b) growing the hES cells *in vitro* in a vessel under conditions under which the cells undergo differentiation and said cells or aggregates thereof do not adhere to the vessel wall:

- (c) incubating for a time sufficient to obtain at least one embryoid body  
5 (EB) containing cells of said defined cell lineage; and  
(d) isolating said cells from said EB.

In accordance with one embodiment, the cells of the defined cell lineage may be inoculated or transplanted into a human recipient to treat a certain human disease or condition, to allow tissue or organ repair, etc. Thus, in accordance with  
10 this preferred embodiment, the cells of the defined cell lineage are formulated in a manner to allow their inoculation or transplantation into the human recipient.

The present invention also provides, by another of its aspects, an inoculable or transplantable preparation comprising cells of said defined cell lineage together with a physiologically acceptable carrier which is compatible with said cells. The  
15 term "*compatible with said cells*" should be understood as a medium which ensures the viability of said cells until they are inoculated or transplanted into the human recipient.

The cells of the defined cell lineage may be induced to undergo further differentiation and transplanted or inoculated into the individual as a cell  
20 suspension or alternatively, they may be cultivated to form a cell mass or an *in vitro* tissue, and the mass or the tissue then being transplanted into the individual.

The hES cells from which the hEBs of the invention are derived may be obtained from established lines (e.g. see Thomson *et al.*, 1998b, *supra*) or may be prepared *de novo* from human embryos which were produced by *in vitro*  
25 fertilization.

## BRIEF DESCRIPTION OF THE DRAWINGS

In order to understand the invention and to see how it may be carried out in practice, a preferred embodiment will now be described, by way of non-limiting  
30 example only, with reference to the accompanying drawings, in which:

Fig. 1A shows cystic hEBs from 3 to 14 days after transfer to petri dishes produced in accordance with the invention. I: simple EB; II & III: cavitated EB; IV: cystic hEB. Scale bar. 40  $\mu$ m.

Fig. 1B shows 5  $\mu$ m paraffin embedded sections of hEBs stained with hematoxylin and eosin (H&E, I,II and IV) or DAPI (III) I: scale bar 80  $\mu$ m; II& III scale bar 20  $\mu$ m; IV: scale bar 4  $\mu$ m.

Fig. 2 shows the expression of cell specific genes in hES cells and cystic hEBs:

Fig. 3 shows *in situ* hybridization analysis of expression of  $\alpha$ -fetoprotein ( $\alpha$ FP),  $\xi$ -globin ( $\xi$ -Glob), neurofilament 68 kd protein (NF) and  $\alpha$ -cardiac actin (cAct) in 20 day old Ebs (Con indicates control):

Fig. 4A shows a pulsating hEB in a relaxed state. Scale bar, 40  $\mu$ m;

Fig. 4B shows the pulsating hEB of Fig. 4A in a subsequent contracted state. Scale bar, 40  $\mu$ m; and

Fig. 4C shows serial sections from a pulsating hEB including a section stained with H&E, a hybridized section showing  $\alpha$ -cardiac actin expression (cActin), and a control section hybridized with non-specific RNA (Control). Scale bar, 100  $\mu$ m.

## 20 MATERIALS AND METHODS

### Formation of human cystic EBs

Human ES cells (H9 clone 10) were grown on mouse embryo fibroblasts in a culture medium consisting of 80% KnockOut™ DMEM (an optimized Dulbecco's modified Eagle's medium for ES cells, Gibco-BRL), 20% KnockOut™ SR (a serum-free formulation, Gibco-BRL), 1 mM glutamine (Gibco-BRL), 0.1 mM  $\beta$ -mercaptoethanol (Sigma), 1% non-essential amino acids stock (Gibco-BRL), 10<sup>3</sup> units/ml leukemia inhibitor factor (LIF) (Gibco-BRL) and 4 ng/ml basic fibroblast growth factor (bFGF) (Gibco-BRL). Under these conditions most of the cells are kept in an undifferentiated state. To induce formation of hEBs, the ES cells were transferred to plastic petri dishes to prevent their adherence to the dish and

promote their aggregation. The concentration of the cells was about  $10^5$  cells per ml. The hEBs were then cultured in the above culture medium not containing leukemia inhibitor factor or basic fibroblast growth factor.

#### 5 Detection of expression of cell specific genes in cystic hEBs.

Total RNA was extracted from cells as previously described (Chirgwin *et al.*, 1979) and cDNA was synthesized from 1  $\mu$ g of total RNA, using a random hexamer (pd(N)<sub>6</sub>) as primer (Pharmacia Biotech) and M-LMV Reverse Transcriptase (Gibco-BRL). cDNA samples were subject to polymerase chain  
10 reaction (PCR) amplification with specific DNA primers. PCR was performed under linear conditions in order to reflect the original amount of the specific transcript. The PCR primers used and the reaction conditions were:  $\alpha$ -fetoprotein: AGAACCTGTCACAAGCTGTG and GACAGCAAGCTGAGGATGTC- Product: 676 base pairs (bp). 20 cycles at 60°C in 1 mM MgCl<sub>2</sub>;  $\xi$ -globin:  
15 GACTGAGAGGACCATCATTG and TCAGGACAGAGGATACGACC - Product: 397 bp. 25 cycles at 60°C in 1 mM MgCl<sub>2</sub>; GAPDH; AGCCACATCGCTCAGACACCA and GTACTCAGCGGCCAGCATCG - Product: 302 bp. 20 cycles at 60°C in 1 mM MgCl<sub>2</sub>. PCR products were analyzed by Western Blot hybridization (Southern, 1975). Probes were radiolabeled by  
20 random priming (Boehringer Mannheim) using [ $\alpha$ -<sup>32</sup>P]dCTP (3000 ci/mM, NEN - Life Science Products).

#### *In situ* hybridization analysis of hEBs

EBs were stained with hematoxylin and eosin (H&E) or hybridized to  
25 specific RNA probes labeled with a fluorescent reagent (Grifman *et al.*, 1998b) and 5  $\mu$ m paraffin embedded serial sections were obtained. Control hybridizations were performed with non-specific RNA. The probes used were 50-mer 2'-O-methyl 5'-biotinylated cRNA of either  $\alpha$ -fetoprotein - TTGTCCCTCTTCAGCAAAGC AGACTTCCTGTTCTGTCCTTGGCAGCATT,  $\xi$ -globin - TGATGGTCCTCT  
30 CAGTCTTGGTCAGAGACATGGCGGCAGGGTGGGCAGCT, Neurofilament



68 kd protein - CCTGCGTGCGGATGGACTTGAGGTCGTTGCTGATGGCG  
GCTACCTGGCTC. or  $\alpha$ -cardiac actin - CGGTGGACAATGGATGGGCCTG  
CCTCATCGTACTCTTGCTTGCTAATCCA.

## 5 EXAMPLES

### Example 1: Formation of cystic hEBs

Fig. 1A shows several hEBs 3 to 14 days after having been transferred to plastic petri dishes and cultured in medium not containing leukemia inhibitor factor or basic fibroblast growth factor. Initially, simple hEBs form (Fig. 1A-I).  
10 Subsequently, the center of the bodies became cavitated (Fig. 1A. II and III), and the bodies began to accumulate fluid, turning to cystic hEBs (Fig. 1A, IV). 20 days after initiation of cellular aggregation most of the structures were cystic and they included a variety of epithelial and mesenchymal cells. Fig. 1B shows 5  $\mu$ m paraffin embedded sections of 20 day old hEBs stained with H&E (I-IV) or DAPI  
15 (V). DAPI staining of the nuclei revealed in some of the cells condensed chromatin that probably corresponds to the apoptosis occurring in the center of the hEB (Fig. 1B-III).

### Example 2: Expression of cell specific genes in cystic hEBs.

20 In order to examine the differentiation status of the EBs, RNA was extracted from hES cells grown on mouse embryo fibroblasts, and from 20 day old EBs. cDNA was synthesized using these RNA samples and expression analysis of several genes was performed by RT-PCR with various human specific DNA primers. The RT-PCR analysis was performed under non-saturating linear  
25 conditions. The identity of the amplified DNA product in the PCR assay was verified by sequence analysis. The lane marked EB in Fig. 2 shows that the hEBs express  $\alpha$ -fetoprotein ( $\alpha$ FP), an endodermal marker (Krumlauf *et al.*, 1985), and  $\xi$ -globin, a marker of hematopoietic cells (Leder *et al.*, 1985)). Very low levels of these markers were observed in the hES cells (Lane ES). The house keeping gene,  
30 glyceraldehyde-3-phosphate dehydrogenase (GAPDH), served as an internal control. Linearity of the observed signal is demonstrated by the "1/10 column"

showing a PCR assay performed with one-tenth the amount of cDNA used in the PCR assay of the hEB sample.

### Example 3: *In situ* hybridization analysis of hEBs

5 To regionally characterize the differentiating cells within the EBs by *in situ* hybridization, expression of four cell-specific molecular markers, all of which transcribe very early during embryonic differentiation, was examined. Serial sections of the hEBs were hybridized with RNA probes specific to  $\alpha$ -fetoprotein ( $\alpha$ FP) (Krumlauf *et al.*, 1985),  $\xi$ -globin ( $\xi$ -glob) ((Leder I., 1985),  $\alpha$ -cardiac actin  
10 (cAct) (Sassoon *et al.*, 1988)) or neurofilament 68kd protein (NF) (Julien *et al.*, 1986)). As shown in Fig. 3, each of these molecular probes specifically labels distinct regions in the EBs. This suggests that the labeled cells in each section were either clonal (derived from a common progenitor cell), or were affected by the same signals and differentiated into the same specific lineage.

15

### Example 4: Characterization of a pulsating hEB

Differentiation of murine ES cells in EBs into the myocardial lineage has been shown to produce pulsating muscle (Sanchez *et al.*, 1991). The existence of cardiac muscle cells in hEBs was demonstrated by *in situ* hybridization of sections  
20 from a human EB with  $\alpha$ -cardiac actin, a marker of embryonic myocardial cells (Sassoon *et al.*, 1988). Figs. 4A and B shows a large vacuolated hEB including cardiac muscle cells that were pulsating at a rate of about 30 beats/min. The hEB was first photographed in a relaxed state (Fig. 4A) and then in a subsequent a contracted state (Fig. 4B). Fig. 4C shows serial sections of a cell including a section  
25 stained with hematoxylin and eosin (H&E), a hybridized section showing  $\alpha$ -cardiac actin expression (cActin), and a section hybridized with non-specific RNA (Control).

## References

- Bain, G., Kitchens, D., Yao, M., Huettner, J.E., Gottlieb, D.I., *Developmental Biology*, **168**:342-57 (1995).
- 5 Capecchi, M.R., *Science*, **244**:1288-92 (1989).
- Chirgwin, J.M., Przybyla, A.E., Macdonald, R.J., Rutter, W.J., *Biochemistry*, **18**:5294-9 (1979).
- 10 Dushnik-Levinson, M., Benvenisty, N., *Biology of the Neonate*, **67**:77-83, (1995).
- Julien, J.P., Meyer, D., Flavell, D., Hurst, J., Grosveld, *Brain Research*, **387**:243-50 (1986).
- 15 Krumlauf, R., Hammer, R.E., Tilghman, S.M., Brinster, R.L., *Molecular and Cellular Biology*, **5**:1639-48 (1985).
- Leder, A., Weir, L., Leder, P., *Molecular and Cellular Biology*, **5**:1025-33 (1985).
- 20 Levinson-Dushnik, M., Benvenisty, N., *Molecular and Cellular Biology*, **17**:3817-22 (1997).
- Lindenbaum, M.H., Grosveld, F., *Genes and Development*, **4**:2075-85 (1990).
- 25 Robertson, E., in *Teratocarcinoma and Embryonic Stem Cells: A Practical Approach*, R. E.J. Eds. (IRL Press, Oxford, 1987) pp. 71-112.
- Rossant, A.L. Joyner, *Trends in Genetics*, **5**:277-83 (1989).
- 30 Sanchez, A., Jones, W.K., Gulick, J., Doetschman, T., Robbins, J., *Journal of Biological Chemistry*, **266**:22419-26 (1991).
- Sassoon, D.A., Garner, I., Buckingham, M., *Development*, **104**:155-64 (1988).
- 35 Southern, E.M., *Journal of Molecular Biology*, **98**:503-17 (1975).
- Thomson, J.A., *et al.*, *Biology of Reproduction*, **55**:254-9 (1996).
- 40 Thomson, J.A., *et al.*, *Proceedings of the National Academy of Sciences of The United States of America*, **92**:7844-8 (1995).
- Thomson, J.A. and Marshall, V.S., *Curr. Top. Dev. Biol.* **38**, 133 (1988a)
- 45 Thomson, J.A., *et al.*, *Science*, **282**:1145-7 (1998b).

Wiles. M.V., Keller. G., *Development*, **111**:259-67 (1991).

Wobus. A.M., Holzhausen. H., Jäkel. P., Schöneich. J., *Experimental Cell*  
5 *Research*, **152**:212-9 (1984).

## CLAIMS:

1. A human embryoid body (hEB).
2. An hEB according to Claim 1, obtained by *in vitro* culturing of human embryonic stem (hES) cells in a vessel under conditions in which the cells undergo  
5 differentiation and the cells or aggregates thereof do not attach to the vessel walls.
3. An hEB according to Claim 1, comprising mesoderm, ectoderm and endoderm lineage cells.
4. An hEB according to Claim 3, comprising cells displaying characteristics of cardiac cells.
- 10 5. An hEB according to Claim 1, comprising cells expressing a marker selected from the group consisting of  $\alpha$ -fetaprotein,  $\xi$ -globin,  $\alpha$ -cardiac actin and neurofilament 68 kd.
6. A process for obtaining at least one human-derived embryoid body (hEB), comprising:
  - 15 (a) providing human embryonic stem (hES) cells;
  - (b) growing the hES cells *in vitro* in a liquid growth medium in a vessel under conditions under which said cells undergo differentiation and said cells or aggregates thereof do not adhere to the vessel wall; and
  - (c) incubating for a time sufficient to develop hEBs from said cells.
- 20 7. A process according to Claim 6, wherein said conditions comprises the absence of leukemia inhibitory factor or fibroblast growth factor from the growth medium.
8. A process according to Claim 6, wherein said condition comprises culturing cells in a vessel having walls of a kind to which cells do not adhere.
- 25 9. A process for preparing cells of a defined cell lineage, comprising:
  - (a) providing human embryonic stem (hES) cells;
  - (b) growing the hES cells *in vitro* in a vessel under conditions in which said cells undergo differentiation and said cells or aggregates thereof do not adhere to the vessel wall;

(c) incubating for a time sufficient to obtain at least one human embryoid body (hEB) containing cells of said defined cell lineage; and

(d) isolating said cells from said hEB.

10. A process according to Claim 9, wherein said cells are formulated for  
5 inoculation or implantation into a recipient individual.


11. An injectable or implantable preparation, comprising cells obtained by the process of Claim 9 or 10 together with a physiologically acceptable carrier compatible with said cells.

12. An implantable cell mass or tissue obtained by the process of Claim 9 or 10.

10 13. Cells produced according to the process according to Claim 9.



**A**



**B**




fig 1

BEST AVAILABLE COPY




fig. 2

BEST AVAILABLE COPY







fig. 3

BEST AVAILABLE COPY




fig. 4

BEST AVAILABLE COPY



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
23 November 2000 (23.11.2000)

PCT

(10) International Publication Number  
**WO 00/70021 A3**

- (51) International Patent Classification<sup>7</sup>: C12N 5/08, (74) Agent: REINHOLD COHN AND PARTNERS; P.O. Box 4060, 61040 Tel Aviv (IL).  
A61K 35/54
- (21) International Application Number: PCT/IL00/00270 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 12 May 2000 (12.05.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 129966 14 May 1999 (14.05.1999) IL (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): YISSUM RESEARCH DEVELOPMENT COMPANY OF THE HEBREW UNIVERSITY OF JERUSALEM [IL/IL]; P.O. Box 4279, 91042 Jerusalem (IL).
- Published:  
— With international search report.
- (71) Applicant and  
(72) Inventor: ITSKOVITZ-ELDOR, Joseph [IL/IL]; Sherit Haplayta St. 42, Danya B., 34987 Haifa (IL).
- (88) Date of publication of the international search report:  
26 April 2001
- (72) Inventor; and  
(75) Inventor/Applicant (*for US only*): BENVENISTY, Nissim [IL/IL]; Hertzl St. 83, Beit-Hakerem, 96344 Jerusalem (IL).
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DIFFERENTIATED HUMAN EMBRYOID CELLS AND A METHOD FOR PRODUCING THEM

(57) Abstract: A process for obtaining human derived embryoid bodies (hEB). Human embryonic stem cells are incubated *in vitro* in a liquid growth medium under conditions in which the cells undergo differentiation, but do not adhere to the walls of the container. The invention also provides hEBs obtained by the process.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/IL 00/00270

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N5/08 A61K35/54

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, MEDLINE, LIFESCIENCES, CHEM ABS Data, EMBASE, SCISEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| A          | THOMSON J A ET AL: "Embryonic stem cell lines derived from human blastocytes." SCIENCE, vol. 282, no. 5391, 6 November 1998 (1998-11-06), pages 1145-1147, XP002151774 cited in the application                       |                       |
| A          | RATHJEN P D ET AL: "Properties and uses of embryonic stem cells: Prospects for application to human biology and gene therapy" REPRODUCTION, FERTILITY AND DEVELOPMENT, vol. 10, no. 1, 1998, pages 31-47, XP000916997 |                       |

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

1 November 2000

Date of mailing of the international search report

15/11/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Teyssier, B


## INTERNATIONAL SEARCH REPORT

Inter. Patent Application No  
PCT/IL 00/00270

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
|------------|--|-----------------------|

|     |   |  |
|-----|---|--|
| P,A | PERA M F ET AL: "Human embryonic stem cells"<br>JOURNAL OF CELL SCIENCES,<br>vol. 113, January 2000 (2000-01), pages 5-10, XP000882172<br>--- |  |
|-----|---|--|



|   |  |   |
|---|--|---|
| A | NAKAMURA T ET AL: "A selective switch-on system for self-renewal of embryonic stem cells using chimeric cytokine receptors."<br>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS,<br>vol. 248, no. 1, 9 July 1998 (1998-07-09), pages 22-27, XP002151775<br>the whole document<br>--- | 7 |
|---|--|---|

|   |   |   |
|---|---|---|
| A | MUMMERY C L ET AL: "Fibroblast growth factor-mediated growth regulation and receptor expression in embryonal carcinoma and embryonic stem cells and human germ cell tumours."<br>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS,<br>vol. 191, no. 1, 1993, pages 188-195, XP002151776<br>the whole document<br>--- | 7 |
|---|---|---|

|     |   |      |
|-----|---|------|
| P,X | ITSKOVITZ-ELDOR J ET AL: "Differentiation of human embryonic stem cells into embryoid bodies comprising the three embryonic germ layers."<br>MOLECULAR MEDICINE,<br>vol. 6, no. 2, February 2000 (2000-02), pages 88-95, XP000953284<br>the whole document<br>----- | 1-13 |
|-----|---|------|



**THIS PAGE BLANK (USPTO)**